

References to list of *Orbiliomycetes* sequences included in this study (Tab. S5)

- Addison, S.L.; Webb, K.; Smaill, S.J.; Menkis, A. – Forest site drives diversity and composition of ectomycorrhizal communities associated with *Pinus radiata*. Unpublished
- Ahrén, D.; Ursing, B.M.; Tunlid, A. (1998). Phylogeny of nematode-trapping fungi based on 18S rDNA sequences. – FEMS Microbiol. Lett. 158: 179–184.
- Amend, A.S.; Seifert, K.A.; Bruns, T.D. (2010). Quantifying microbial communities with 454 pyrosequencing: does read abundance count? – Mol. Ecol. 19: 5555–5565.
- Arenz, B.E. & Blanchette, R.A. (2009). Investigations of fungal diversity in wooden structures and soils at historic sites on the Antarctic Peninsula. – Can. J. Microbiol. 55: 46–56.
- Arhipova, N.; Gaitnieks, T.; Donis, J.; Stenlid, J.; Vasaitis, R. (2011). Decay, yield loss and associate fungi in stands of grey alder (*Alnus incana*) in Latvia. – Forestry 84: 337–348.
- Avis, P.G. & Norris, M. – Fungi colonizing litter decomposition bags at Cedar Creek Ecosystem Science Reserve, Minnesota, USA. Unpublished
- Baier, C. – Molecular analysis of ecologically important fungi. Unpublished
- Ban, S.; Mayuzumi, S.; Tabuchi, Y.; Shimamura, K.; Inaba, S. – Unpublished
- Beck, A.M. & Yannarell, A.C. – Unpublished
- Beck, A.M.; Dawson, J.O.; Yannarell, A.C. – Linkages between potential root-bound symbionts and non-symbiotic soil microbes in the alder and birch rhizosphere. Unpublished
- Belliveau MJR, Bärlocher F (2005). Molecular evidence confirms multiple origins of aquatic hyphomycetes. – Mycol. Res. 109: 1407–1417.
- Berdoulay, M. & Salvado, J.C. (2009). Genetic characterization of microbial communities living at the surface of building stones. – Appl. Microbiol. 49: 311–316.
- Bhilabutra, W. (2009). Biodiversity of fungi on some grasses and their production of bioactive compounds. Chapter 5: Ribosomal DNA phylogenies reveal that *Dactylaria* is polyphyletic. – Thesis, Chiang Mai University, Thailand. http://archive.lib.cmu.ac.th/full/T/2009/biol0209wb_ch5.pdf
- Brown, S.P.; Ferrer, A.; Dalling, J.W.; Heath, K.D. (2016). Don't put all your eggs in one basket: a cost-effective and powerful method to optimize primer choice for rRNA environmental community analyses using the Fluidigm Access Array. – Mol. Ecol. Resour. 16: 946–956.
- Campos-Herrera, R. – Unpublished
- Castaneda-Ramirez, G.S.; Mendoza-de-Gives, P.; Aguilar-Marcelino, L.; Lopez-Arellano, M.E.; Hernandez-Romano, J. (2016). Phylogenetic analysis of nucleotide sequences from the ITS region and biological characterization of nematophagous fungi from Morelos, Mexico. – J Mycology 2016, Article ID 8502629, 13 pages.
- Castillo, B.T.; LeMoine, J.; Nave, L.; James, T.Y.; Nadelhoffer, K.J. – Impacts of experimentally accelerated forest succession on belowground plant and fungal communities. Unpublished
- CBS (<http://www.westerdijkinstituut.nl/Collections/BioloMICSSequences.aspx>)
- Chen, C. Y. & Liu, P. W. – Records of fungi in Taiwan. Unpublished
- Chen, J.; Xu, L.L.; Liu, B.; Liu, X.Z. (2007). Taxonomy of *Dactylella* complex and *Vermispora*. I. Generic concepts based on morphology and ITS sequences data. – Fungal Diversity 26: 73–83.
- Cox, F.; Newsham, K.K.; Bol, R.; Dungait, J.A.; Robinson, C.H. (2016). Not poles apart: Antarctic soil fungal communities show similarities to those of the distant Arctic. – Ecol. Lett. 19: 528–536.
- Crous, P.W. (2010). *Amphospora persooniae*. Fungal Planet 669. – Persoonia 39: 370–371.
- Curlevski, N.J.A.; Cairney, J.W.G.; Anderson, I.C.; Xu, Z.H. – Soil fungal communities differ in native mixed forest and adjacent *Araucaria cunninghamii* plantations in subtropical Australia. Unpublished
- Edwards, I.P. & Zak, D.R. (2011). Fungal community composition and function after long-term exposure of northern forests to elevated atmospheric CO₂ and tropospheric O₃. – Glob Chang Biol 17: 2184–2195.
- Eichorst, S.A. & Kuske, C.R. (2012). Identification of cellulose-responsive bacterial and fungal communities in geographically and edaphically different soils by using stable isotope probing. – Appl. Environ. Microbiol. 78: 2316–2327.
- Ekanayaka, A.H.; Hyde, K.D.; Jones, E.B.G.; Zhao, Q. (2018). *Orbiliaceae* from Thailand. – Mycosphere 9: 155–168.
- Findley, K.; Oh, J.; Yang, J.; Conlan, S.; Deming, C.; Meyer, J.A.; Schoenfeld, D.; Nomicos, E.; Park, M.; NIH Intramural Sequencing Center Comparative Sequencing Program; Kong, H.H.; Segre, J.A. (2013). Topographic diversity of fungal and bacterial communities in human skin. Nature 498: 367–370.
- Fröhlich-Nowoisky, J.; Despres, V.R.; Pöschl, U. – Global atmospheric diversity of fungi: *Asco*- and *Basidiomycota* in continental and marine air. Unpublished
- Gao, Q. & Yang, Z.L. – Ectomycorrhizal fungi associated with *Polygonum* and *Potentilla* species on an alpine-meadow in southwestern China. Unpublished
- Gao, C.; Shi, N.N.; Chen, L.; Ji, N.N.; Wu, B.W.; Wang, Y.L.; Xu, Y.; Zheng, Y.; Mi, X.C.; Ma, K.P.; Guo, L. (2017). Relationships between soil fungal and woody plant assemblages differ between ridge and valley habitats in a subtropical mountain forest. – New Phytologist 213: 1874–1885.
- Glinka, C.B. & Hawkes, C.V. (2014). Environmental controls on fungal community composition and abundance over three years in native and degraded shrublands. – Microbial Ecology 68: 807–817.

- Gonzalez-Chavez, M.C.C.; Torres-Cruz, T.J.; Albarran-Sanchez, S.; Porras-Alfaro, A.; Carrillo-Gonzalez, R.; Carrillo-Lopez, L.M. – Fungi involved in peloton colonization in *Vanilla* species from different cultivation systems. Unpublished
- Gouveia, G.V.; Yano de Melo, A.M.; da Costa, M.M.; Gouveia, J. – Soil microbial diversity of a Brazilian semiarid region. Unpublished
- Grau, O.; Geml, J.; Perez-Haase, A.; Ninot, J.M.; Semenova-Nelsen, T.A.; Penuelas, J. (2017). Abrupt changes in the composition and function of fungal communities along an environmental gradient in the high Arctic. – *Mol. Ecol.* 26: 4798–4810.
- Guichon, S.H.A. & Simard, S.W. – Diversity of fungi found in soils from Cedar-Hemlock forests and Hemlock-Amabilis fir forests. Unpublished
- Guo, J.W.; Yu, Z.F.; Qiao, M.; Zhang, K.Q. (2009). Effects of carbon and nitrogen sources on sexual reproduction of five strains from the ascomycete *Orbilia*. – *Ann. Microbiol.* 59: 1–5.
- Haelewaters, D.; Dirks, A.C.; Kappler, L.A.; Mitchell, J.K.; Quijada, L.; Vandegrift, R.; Buyck, B.; Pfister, D.H. (2018). A preliminary checklist of fungi at the Boston Harbor Islands. *Northeastern Naturalist* 25 (sp9): 45–76.
- Hagedorn, G.; Scholler, M. (1999). A reevaluation of predatory orbiliaceous fungi. I. Phylogenetic analysis using rDNA sequence data. – *Sydowia* 51: 27–48.
- Han, Z.G.; Sui, X.; Li, M.S. (2016). Effects of forest age on soil fungal community in a northern temperate ecosystem. *Indian J. Microbiol.* 56: 328–334.
- Harrington, F.A.; Pfister, D.H.; Potter, D.; Donoghue, M. (1999). Phylogenetic studies within the *Pezizales*: I. 18S rRNA sequence data and classification. – *Mycologia* 91: 41–50.
- Hartmann, M.; Lee, S.; Hallamand, S.J.; Mohn, W.W. (2009). Bacterial, archaeal and eukaryal community structures throughout soil horizons of harvested and naturally disturbed forest stand. – *Environm. Microbiol.* 11: 3045–3062.
- Hassett, J.E. & Zak, D.R. – Nitrogen deposition alters fungal community composition in northern hardwood forests. Unpublished
- Hawkes, C.V.; Kivlin, S.N.; Rocca, J.D.; Huguet, V.; Thomsen, M.; Suttle, K.B. (2011). Fungal community responses to precipitation. *Global Change Biology* 17: 1637–1645.
- Held, B.W. & Blanchette, R.A. (2017). Deception Island, Antarctica, harbors a diverse assemblage of wood decay fungi. *Fungal Biol.* 121: 145–157.
- Hernández-Restrepo, M.; Groenewald, J.Z.; Crous, P.W. (2015) [2016]. Taxonomic and phylogenetic re-evaluation of *Microdochium*, *Monographella* and *Idriella*. – *Persoonia* 36: 57–82.
- Herrera, J.; Khidir, H.H.; Eudy, D.M.; Porras-Alfaro, A.; Natvig, D.O.; Sinsabaugh, R.L. (2010). Shifting fungal endophyte communities colonize *Bouteloua gracilis*: effect of host tissue and geographical distribution. – *Mycologia* 102: 1012–1026.
- Herrmann, A.; Kainiemi, V.; Rosling, A.; Urbina, H.; Boye, K.; Fransson, P. – Linking bioenergetics to microbial communities along a Scots Pine stand age gradient. Unpublished
- Hollister, E.B.; Schadt, C.W.; Palumbo, A.V.; Ansley, R.J.; Boutton, T.W. (2010). Structural and functional diversity of soil bacterial and fungal communities following woody plant encroachment in the southern Great Plains. – *Soil Biol. Biochem.* 42: 1816–1824.
- Houseknecht, J.L. & Zhou, J. – Unpublished
- Jakubowicz, I.; Yarahmadi, N.; Arthurson, V. (2011). Kinetics of abiotic and biotic degradability of low-density polyethylene containing prodegradant additives and its effect on the growth of microbial communities. – *Polymer degradation and Stability* 96: 919–928.
- James, T.Y.; Letcher, P.M.; Longcore, J.E.; Mozley-Standridge, S.E.; Porter, D.; Powell, M.J.; Griffith, G.W.; Vilgalys, R. (2006). A molecular phylogeny of the flagellated fungi (*Chytridiomycota*) and description of a new phylum (*Blastocladiomycota*). – *Mycologia* 98: 860–871.
- James, T.Y.; Marino, J.A.; Perfecto, I.; Vandermeer, J. (2016). Identification of putative coffee rust mycoparasites via single-molecule DNA sequencing of infected pustules. – *Appl Environ Microbiol* 82: 631–639.
- Ji, K.Y.; Yu, Z.F.; Zhang, Y.; Zhang, K.Q. – Unpublished
- Jiang, X.; Liu, B.; Liu, X.Z. – *Orbilia* species and their anamorphs. Unpublished
- Jumpponen, A.; Jones, K.L.; Blair, J. (2010). Vertical distribution of fungal communities in tallgrass prairie soil. – *Mycologia* 102: 1027–1041.
- Kaminsky, L.; Smith, M.E.; Healy, R.; Grupe, A.; Spakes-Richter, B. – Fungi of the Ordway Swisher Biological Station. Unpublished
- Kano, S.; Aimi, T.; Masumoto, S.; Kitamoto, Y.; Morinaga, T. (2004). Physiology and molecular characteristics of a pine wilt nematode-trapping fungus, *Monacrosporium megalosporum*. – *Curr. Microbiol.* 49: 158–164.
- Karst, J.; Piculell, B.; Brigham, C.; Booth, M.; Hoeksema, J.D. (2013). Fungal communities in soils along a vegetative ecotone. – *Mycologia* 105: 61–70.
- Kauserud, H.; Mathiesen, C.; M. Ohlson (2008). High diversity of fungi associated with living parts of boreal forest bryophytes. *Botany* 86: 1326–1333.
- Kim, D.G.; Ryu, Y.H.; Hwang, H.G. (2006). First report of two nematode-trapping fungi, *Monacrosporium ullum* sp. nov. and *Arthrobotrys amerospora*, from Korea. – *Plant Pathol. J.* 22: 174–178.
- Kivlin, S.N. & Hawkes, C.V. (2011). Differentiating between effects of invasion and diversity: impacts of aboveground plant communities on belowground fungal communities. – *New Phytol.* 189: 526–535.
- Kubart, A.; Vasaitis, R.; Stenlid, J.; Dahlberg, A. (2016). Fungal communities in Norway spruce stumps along a longitudinal gradient in Sweden. – *Forest Ecology and Management* 371: 10.1016/j.foreco.2015.12.017.

- Kuhnert, R.; Oberkofler, I.; Peintner, U. (2012). Fungal growth and biomass development is boosted by plants in snow-covered soil. – *Microbial Ecology* 64: 79–90.
- Kumar, T.K.A.; Healy, R.; Spatafora, J.W.; Blackwell, M.; McLaughlin, D.J. (2012). *Orbilia* ultrastructure, character evolution and phylogeny of *Pezizomycotina*. – *Mycologia* 104: 462–476.
- Kuo, C.C.; Chen, P.; Chen, C.Y. (2008). New records of nematode-trapping fungi from Taiwan. – *Fung. Sci.* 23: 27–34.
- Lazarevic & Menkis 2020
- Lesaulnier, C.; Papamichail, D.; McCorkle, S.; Ollivier, B.; Skiena, S.; Taghavi, S.; Zak, D.; van der Lelie, D. (2008). Elevated atmospheric CO₂ affects soil microbial diversity associated with trembling aspen. – *Environ. Microbiol.* 10: 926–941.
- Lee, R.W. – Comparative metagenomics reveals herpotrichiellaceous core in diverse rhizome mycobiomes inhabiting different mycoheterotrophic *Gastrodia* orchids. Unpublished
- Li, J.Y. – Two new anamorphic species of *Dactylella* sp. nov. from China. Unpublished
- Li, C. & Liang, J. – High diversity of the soil fungi from Yangdongshan Shierdushui Nature Reserve. Unpublished
- Li, Y.; Hyde, K.D.; Jeewon, R.; Cai, L.; Vijaykrishna, D.; Zhang, K.Q. (2005). Phylogenetics and evolution of nematode-trapping fungi (*Orbiliaceae*) estimated from nuclear and protein coding genes. – *Mycologia* 97: 1034–1046.
- Li, Y.; Jeewon, R.; Hyde, K.D.; Mo, M.H.; Zhang, K.Q. (2006). Two new species of nematode-trapping fungi: relationships inferred from morphology, rDNA and protein gene sequence analyses. – *Mycol. Res.* 110: 790–800.
- Li, J.; Liu, Y.; Zhu, H.Y.; Zhang, K.Q. (2016). Phylogenetic analysis of adhesion related genes Mad1 revealed a positive selection for the evolution of trapping devices of nematode-trapping fungi. – *Scientific Reports* 6: 22609. doi: 10.1038/srep22609.
- Li, J.Y.; Qian, W.Y.; Qiao, M.; Bai, Y.L.; Yu, Z.F. (2013). A new *Drechlerella* species from Hainan, China. – *Mycotaxon* 125: 183–188.
- Li, J.; Yang, J.K.; Liang, L.M.; Zhang, K.Q. (2008). Taxonomic revision of the nematode-trapping fungus *Arthrobotrys multisecondaria*. – *J. Microbiol.* 46: 513–518.
- Li, S.F.; Yu, Z.F.; Zhang, Y.; Qiao, M.; Guo, J.W.; Zhang, K.Q. (2009). *Arthrobotrys nonseptata*, a new anamorph from an *Orbilia* species. – *Mycotaxon* 109: 247–254.
- Li, W.P.; Zhang, K.Q.; Xing, L. J.; Liew, E.C.; Hyde, K.D. – Systematic state of *Arthrobotrys guizhouensis* inferred from large subunit ribosomal DNA sequence. Unpublished
- Li, W.P.; Zhang, K.Q.; Xing, L.J.; Liew, E.C.Y.; Hyde, K.D. – Phylogenetic analysis of predatory anamorphic *Orbiliaceae* based on sequences of 28S large subunit ribosomal RNA gene and a new generic concept. Unpublished
- Lindner, D.L.; Vasaitis, R.; Kubartova, A.; Allmer, J.; Johannesson, H.; Banik, M.T.; Stenlid, J. (2011). Initial fungal colonizer affects mass loss and fungal community development in *Picea abies* logs 6 yr after inoculation. – *Fungal Ecol* 4: 449–460.
- Lipson, D.A.; Kuske, C.R.; Gallegos-Graves, L.V.; Oechel, W.C. (2014). Elevated atmospheric CO₂ stimulates soil fungal diversity through increased fine root production in a semiarid shrubland ecosystem. – *Glob. Chang. Biol.* 20: 2555–2565.
- Liou, G.Y.; Tzean, S.S. (1997). Phylogeny of the genus *Arthrobotrys* and allied nematode-trapping fungi based on rDNA sequences. – *Mycologia* 89: 876–884.
- Liu, P.W. & Chen, C.Y. – Records of fungi in Taiwan. Unpublished
- Liu, B.; Liu, X.Z.; Zhuang, W.Y. (2005a). *Orbilia querci*, sp. nov. and its knob-forming nematophagous anamorph. – *FEMS Microbiol. Lett.* 245: 99–105.
- Liu, B.; Liu, X.Z.; Zhuang, W.Y. (2005b). A new species of *Hyalorbilia* and its anamorph from China. – *Nova Hedwigia* 81: 145–155.
- Liu, B.; Liu, X.Z.; Zhuang, W.Y. (2006). Taxonomy and molecular phylogeny of *Orbiliaceae* from China. – Poster presented at the 8th International Mycological Congress in August 2006 in Cairns, Australia.
- Liu, S.H.; Su, H.Y.; Su, X.J.; Zhang, F.; Liao, G.G.; Yang, X.J. (2014). *Arthrobotrys xiangyunensis*, a novel nematode-trapping taxon from a hot-spring in Yunnan Province, China. *Phytotaxa* 174: 89–96.
- Liu, X.; Yang, Y.; Sun, M. – A new variety of nematode-trapping fungi forming adhesive hyphal branch. Unpublished
- Liu, X.; Zhang, W.; Liu, K. – *Drechlerella stenobrocha* genome reveals carnivorous origination and mechanical trapping mechanism of predatory fungi. Unpublished
- Macia-Vicente, J.G.; Ferraro, V.; Burruano, S.; Lopez-Llorca, L.V. (2012). Fungal assemblages associated with roots of halophytic and non-halophytic plant species vary differentially along a salinity gradient. – *Microb. Ecol.* 64: 668–679.
- Magyar, D.; Merényi, Z.; Bratek, Z.; Baral, H.O.; Marson, G. (2016). *Lecophagus vermicola* sp. nov., a nematophagous hyphomycete with an unusual hunting strategy. – *Mycol. Progr.* 15: 1137–1144.
- Magyar, D.; Merényi, Z.; Körmöczy, P.; Bratek, Z.; Kredics, L. (2017a). Phylogenetic analysis and description of two new species of pollen-parasitic *Retarius* (anamorphic *Orbiliomycetes*). – *Nova Hedwigia* 105: 411–423.
- Magyar, D.; Merényi, Z.; Udvardy, O.; Kajtor-Apatini, D.; Körmöczy, P.; Fülöp, A.; Bratek, Z.; Kredics, L. (2017b)[2018]. *Mycoceros antenatissimus* gen. et sp. nov.: a mitosporic fungus capturing pollen grains. – *Mycol. Progr.* 17: 33–43.
- Martos, F.; Munoz, F.; Pailler, T.; Kottke, I.; Gonneau, C.; Selosse, M.A. (2012). The role of epiphytism in architecture and evolutionary constraint within mycorrhizal networks of tropical orchids. – *Mol. Ecol.* 21, 5098–5109.
- McGuire, K.L.; Zak, D.R.; Edwards, I.P.; Blackwood, C.B.; Upchurch, R. (2010). Slowed decomposition is biotically mediated in an ectomycorrhizal, tropical rain forest. – *Oecologia* 164, 785–795.

- Meerupati, T.; Andersson, K.M.; Friman, E.; Kumar, D.; Tunlid, A.; Ahren, D. (2013). Genomic mechanisms accounting for the adaptation to parasitism in nematode-trapping fungi. – *PLoS Genet.* 9: E1003909
- Menkis, A.; Vasiliauskas, R.; Taylor, A.F.S.; Stenström, E.; Stenlid, J.; Finlay, R. (2006). Fungi in decayed roots of conifer seedlings in forest nurseries, afforested clear-cuts and abandoned farmland. – *Plant Pathol.* 55: 117–129.
- Menkis, A.; Burokienė, D.; Gaitnieks, T.; Uotila, A.; Johannesson, H.; Rosling, A.; Finlay, R.D.; Stenlid, J.; Vasaitis, R. (2012). Occurrence and impact of the root-rot biocontrol agent *Phlebiopsis gigantea* on soil fungal communities in *Picea abies* forests of northern Europe. – *FEMS Microbiol. Ecol.* 81: 438–445.
- Menkis, A.; Marčiulynas, A.; Gedminas, A.; Lynikienė, J.; Povilaitienė, A. (2015). High-throughput sequencing reveals drastic changes in fungal communities in the phyllosphere of Norway spruce (*Picea abies*) following invasion of the spruce bud scale (*Physokermes piceae*). – *Microb. Evol.* 70: 904–911.
- Menkis, A.; Östbrant, I.L.; Davydenko, K.; Bakys, R.; Balalaikins, M.; Vasaitis, R. (2016). *Scolytus multistriatus* associated with Dutch elm disease on the island of Gotland: phenology and communities of vectored fungi. – *Mycol. Progr.* 15:55 p. 1–8. doi:10.1007/s11557-016-1199-3
- Meyer, S.L.F.; Carta, L.K.; Rehner, S.A. (2005). Morphological variability and molecular phylogeny of the nematophagous fungus *Monacrosporium drechleri*. – *Mycologia* 97: 405–415.
- Miyazaki, K.; Tsuchiya, Y.; Okuda, T. – Filamentous fungi isolated in Japanese mushroom cultivation facilities. Unpublished
- Mo, M.H.; Huang, X.W.; Zhou, W.; Huang, Y.; Hao, Y.E. (2005). *Arthrobotrys yunnanensis*, sp. nov., the fourth anamorph of *Orbilia auricolor*. – *Fungal Diversity* 1: 107–115.
- Moll, J.; Hoppe, B.; König, S.; Wubet, T.; Buscot, F.; Krüger, D. (2016). Spatial distribution of fungal communities in an arable soil. *PLoS ONE* 11, E0148130
- Mueller, R.C.; Balash, M.M.; Kuske, C.R. (2014). Contrasting soil fungal community responses to experimental nitrogen addition using the large subunit rRNA taxonomic marker and cellobiohydrolase I functional marker. – *Mol Ecol.* 23: 4406–17. doi: 10.1111/mec.12858. Epub 2014 Aug 7.
- Murga-Gutierrez, S.N.; Colagiero, M.; Rosso, L.C.; Finetti Sialer, M.M.; Ciancio, A. (2012). Root-knot nematodes from asparagus and associated biological antagonists in Peru. – *Nematropica* 42: 57–62.
- Nagesh, M.; Saleem, J.; Thippeswamy, R.; Nikhita, P.R.; Balachander, M. – Unpublished
- Napoli, C.; Mello, A.; Borra, A.; Vizzini, A.; Sourzat, P.; Bonfante, P. (2010). *Tuber melanosporum*, when dominant, affects fungal dynamics in truffle grounds. – *New Phytol.* 185: 237–247.
- Nemergut, D.R.; Townsend, A.R.; Sattin, S.R.; Freeman, K.R.; Fierer, N.; Neff, J.C.; Bowman, W.D.; Schadt, C.W.; Weintraub, M.N.; Schmidt, S.K. (2008). The effects of chronic nitrogen fertilization on alpine tundra soil microbial communities: implications for carbon and nitrogen cycling. – *Environ. Microbiol.* 10: 3093–3105.
- Nishizawa, T.; Zhaorigetu, Komatsuzaki, M.; Sato, Y.; Kaneko, N.; Ohta, H. (2010). Molecular characterization of fungal communities in non-tilled, cover-cropped upland rice field soils. – *Microbes Environ.* 25: 204–210.
- Oberkofler, I.; Peintner, U. – Another view of soil mycota: soil washing - an effective method for exclusion of spore banks from soil clone libraries. Unpublished
- Oberkofler, I.; Peintner, U. (2008). Detection of soil fungal communities in an alpine primary successional habitat: Does pooling of DNA extracts affect investigations? – *Ann. Microbiol.* 58: 585–595.
- Ohori, A.; Endo, S.; Sano, A.; Yokoyama, K.; Yarita, K.; Yamaguchi, M.; Kamei, K.; Miyaji, M.; Nishimura, K. (2006). Rapid identification of *Ochroconis gallopava* by a loop-mediated isothermal amplification (LAMP) method. – *Vet. Microbiol.* 114: 359–365.
- Olatinwo, R.; Yin, B.; Becker, J.O.; Borneman, J. – An approach for identifying microorganisms involved in specific functions: detection of fungi that suppress *Heterodera schachtii*. Unpublished
- Parchert, K.J.; Spilde, M.N.; Porras-Alfaro, A.; Nyberg, A.M.; Northup, D.E. (2012). Fungal communities associated with rock varnish in Black Canyon, New Mexico: Casual inhabitants or essential partners? – *Geomicrobiology Journal* 29: 752–766.
- Parrent, J.L. & Vilgalys, R. (2007). Biomass and compositional responses of ectomycorrhizal fungal hyphae to elevated CO₂ and nitrogen fertilization. – *New Phytol.* 176: 164–174.
- Pathak, E.; El-Borai, F.E.; Campos-Herrera, R.; Johnson, E.G.; Stuart, R.J.; Graham, J.H.; Duncan, L.W. (2012). Use of real-time PCR to discriminate parasitic and saprophagous behaviour by nematophagous fungi. – *Fungal Biol* 116: 563–573.
- Pfister, D.H. (1997). Castor, Pollux and life histories of fungi. – *Mycologia* 89: 1–23.
- Pitkaranta, M.; Meklin, T.; Hyvarinen, A.; Paulin, L.; Auvinen, P.; Nevalainen, A.; Rintala, H. (2008). Analysis of fungal flora in indoor dust by ribosomal DNA sequence analysis, quantitative PCR, and culture. – *Appl. Environ. Microbiol.* 74: 233–244.
- Polishook, J.D.; Moncalvo, J.M.; Vilgalys, R.; Bills, G. – Diversity and distributions of saprobic cord-forming basidiomycetes in two small patches of forest litter. Unpublished
- Qiao, M.; Li, J.Y.; Baral, H.O.; Zhang, Y.; Qian, W.Y.; Su, H.Y.; Yu, Z.F. (2015). *Orbilia yuanensis* sp. nov. and its anamorph. – *Mycol. Progr.* 14: 1022: 1–7.
- Qiao, M.; Zhang, Y.; Li, S.F.; Baral, H.O.; Weber, E.; Su, H.Y.; Xu, J.P.; Zhang, K.Q.; Yu, Z.F. (2012). *Orbilia blumenaviensis* and its *Arthrobotrys* anamorph. – *Mycol. Progr.* 11: 255–262.
- Qiao, M.; Zhang, Y.; Baral, H.; Zhang, K.Q.; Yu, Z.F. – New species and new combination of *Orbiliaceae* from Yunnan, China. Unpublished

- Quijada, L.; Baral, H.O.; Jaen-Molinia, R.; Weiß, M.; Caujapé-Castells, J.; Beltrán-Tejera, E. (2014). Phylogenetic and morphological circumscription of the *Orbilia aurantiorubra* group. – *Phytotaxa* 175: 1–18.
- Qin, L.; Qiao, M.; Yang, Y.; Yang, G.Z.; Lu, K.P.; Zhang, K.Q.; Xu, J.P.; Yu, Z.F. (2011). *Dactylella pseudobrevistipitata*, a new species from China. – *Ann. Microbiol.* 61: 591–595.
- Rodrigues, D.F.; Jaisi, D.P.; Elimelech, M. (2013). Toxicity of functionalized single-walled carbon nanotubes on soil microbial communities: implications for nutrient cycling in soil. – *Environ. Sci. Technol.* 47: 625–633.
- Rosales-Castillo, J.A.; Oyama, K.; Vazquez-Garciduenas, M.S.; Aguilar-Romero, R.; Garcia-Oliva, F.; Vazquez-Marrufo, G. (2018). Fungal community and ligninolytic enzyme activities in *Quercus deserticola* Trel. litter from forest fragments with increasing levels of disturbance. – *Forests* 9: 11.
- Rosling, A.; Midgley, M.G.; Cheeke, T.; Urbina, H.; Fransson, P.; Phillips, R.P. (2016). Phosphorus cycling in deciduous forest soil differs between stands dominated by ecto- and arbuscular mycorrhizal trees. – *New Phytol.* 209: 1184–1195.
- Sanchez-Martinez, E. & Mendoza-de-Gives, P. – Unpublished
- Sayer, E.J.; Wagner, M.; Oliver, A.E.; Pywell, R.F.; James, P.; Whiteley, A.S.; Heard, M.S. (2013). Grassland management influences spatial patterns of soil microbial communities. *Soil Biology and Biochemistry* 61: 61–68.
- Schmalenberger, A. & Noll, M. – The effect of sulfonate addition to bacterial and fungal communities in grasslands, conclusions on the microbial availability of carbon bonded sulphur. Unpublished
- Schmidt, S.K. & Darcy, J.L. (2014). Phylogeny of ulotrichalean algae from extreme high altitude and high latitude ecosystems. – *Polar Biol.* 38: 689–697.
- Schoch, C.L.; Sung, G.H.; López-Giráldez, F. et al. (2009). The *Ascomycota* tree of life: a phylum-wide phylogeny clarifies the origin and evolution of fundamental reproductive and ecological traits. – *Syst. Biol.* 58: 224–239.
- Semenova, T.A.; Morgado, L.N.; Welker, J.M.; Walker, M.D.; Smets, E.; Geml, J. (2015). Long-term experimental warming alters community composition of ascomycetes in Alaskan moist and dry arctic tundra. – *Mol. Ecol.* 24, 424–437.
- Shao, Y.Y.; Baral, H.O.; Ou, X.Y.; Wu, H.; Huang, F.C.; Zheng, H.F.; Liu, B. (2018). New species and records of orbiliaceous fungi from Georgia, USA. – *Mycol. Progr.* 17: 1225–1235.
- Skipp, R.A.; Yeates, G.W.; Chen, L.Y.; Glare, T.R. (2002). Occurrence, morphological characteristics and ribotyping of New Zealand isolates of *Duddingtonia flagrans*, a candidate for biocontrol of animal parasitic nematodes. – *New Zealand Journal of Agricultural Research*, 45: 187–196.
- Smith, M.E.; Jaffee, B.A. (2009). PCR primers with enhanced specificity for nematode-trapping fungi (*Orbiliaceae*). – *Microb. Ecol.* 58: 117–128.
- Song, P.; Tanabe, S.; Yi, R.; Kagami, M.; Liu, X.; Ban, S. (2018). Fungal community structure at pelagic and littoral sites in Lake Biwa determined with high-throughput sequencing. – *Limnology* 19: 241–251.
- Spatafora, J.W.; Sung, G.H.; Johnson, D. et al. (2007, '2006'). A five-gene phylogeny of *Pezizomycotina*. – *Mycologia* 98: 1018–1028.
- Steven, B.; Gallegos-Graves, L.V.; Yeager, C.; Belnap, J.; Kuske, C.R. (2013). Common and distinguishing features of the bacterial and fungal communities in biological soil crusts and shrub root zone soils. – *Soil Biol. Biochem.* 69: 302–312.
- Stirling, G.; Rames, E.; Stirling, M.; Hamill, S. (2011). Factors associated with the suppressiveness of sugarcane soils to plant-parasitic nematodes. – *J. Nematol.* 43: 135–148.
- Stone, K.M.E.; Leigh, M.B.; Taylor, D.L.; Valentine, D. – Identification of cellulose utilizing soil microorganisms by stable isotope probing (SIP). Unpublished
- Su, H.Y.; Hao, Y.; Liu, S.R.; Li, Y.X.; Cao, Y.H.; Chen, M.H.; Su, X.J.; Yang, X.Y. (2011a). *Dactylella qiluensis*, a new species of aquatic hyphomycete from China. – *Cryptog. Mycol.* 32: 177–183.
- Su, H.Y.; Liu, S.R.; Li, Y.X.; Cao, Y.H.; Chen, M.H.; Yang, X.Y. (2011b). *Arthrobotrys latispora*, a new nematode-trapping fungus from southwest China. – *Mycotaxon* 117: 29–36.
- Suvi, T.; Tedersoo, L.; Abarenkov, K.; Beaver, K.; Gerlach, J.; Koljalg, U. (2010). Mycorrhizal symbionts of *Pisonia grandis* and *P. sechellarum* in Seychelles: identification of mycorrhizal fungi and description of new *Tomentella* species. – *Mycologia* 102: 522–533.
- Swe, A.; Jeewon, R.; Pointing, S.B.; Hyde, K.D. (2008). Taxonomy and molecular phylogeny of *Arthrobotrys mangrovispora*, a new marine nematode-trapping fungal species. – *Bot. Mar.* 51: 331–338.
- Swe, A., Jeewon, R., Pointing, S.B. and Hyde, K.D. – Ecological speciation and molecular phylogeny of genus *Arthrobotrys* based on multi-gene sequence analyses. Unpublished
- Tanabe, Y.; Nagahama, T.; Saikawa, M.; Sugiyama, J. (1999). Phylogenetic relationship of *Cephalophora* to nematophagous hyphomycetes including taxonomic and nomenclatural emendations of the genus *Lecophagus*. – *Mycologia* 91: 830–835.
- Taylor, D.L.; Hollingsworth, T.N.; McFarland, J.W.; Lennon, N.J.; Nusbaum, C.; Ruess, R.W. (2014). A first comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale niche partitioning. – *Ecological Monographs* 84: 3–20.
- Tedersoo, L.; Bahram, M.; Puusepp, R.; Nilsson, R.H.; James, T.Y. (2017). Novel soil-inhabiting clades fill gaps in the fungal tree of life. – *Microbiome* 5: 42, doi 10.1186/s40168-017-0259-5
- Timling, I.; Walker, D.A.; Nusbaum, C.; Lennon, N.J.; Taylor, D.L. (2014). Rich and cold: diversity, distribution and drivers of fungal communities in patterned-ground ecosystems of the North American Arctic. – *Mol. Ecol.* 23: 3258–3272.
- Thornton, C. – Development of a lateral-flow device for the specific detection of *Trichoderma* species in soil and in the plant rhizosphere. Unpublished

- Unterseher, M.; Persöh, D.; Schnittler, M. (2013). Leaf-inhabiting endophytic fungi of European Beech (*Fagus sylvatica* L.) co-occur in leaf litter but are rare on decaying wood of the same host. – *Fungal Diversity* 60: 43–54.
- Urbina, H.; Scofield, D.G.; Cafaro, M.; Rosling, A. (2016). DNA-metabarcoding uncovers the diversity of soil-inhabiting fungi in the tropical island of Puerto Rico. – *Mycoscience* 57: 217–227.
- Van Agtmaal, M. – Unpublished
- Vu, D.; Groenewald, M.; de Vries, M.; Gehrmann, T.; Stielow, B.; Eberhardt, U.; Al-Hatmi, A.; Groenewald, J.Z.; Cardinali, G.; Houbraken, J.; Boekhout, T.; Crous, P.W.; Robert, V.; Verkley, G.J.M. (2019). Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. – *Stud. Mycol.* 92: 135–154.
- Waldrop, M.P.; Zak, D.R.; Blackwood, C.B.; Curtis, C.D.; Tilman, D. (2006). Resource availability controls fungal diversity across a plant diversity gradient. – *Ecol. Lett.* 9: 1127–1135.
- Wang, J.; Meng, Q.; Qiao, J.; Wang, W. – Cloning and sequence analysis of ribosomal RNA genes of *Arthrobotrys oligospora* strain XAol1. Unpublished
- Wang, Z.; Wang, C.Y.; Gu, L.J.; Sun, B.S.; Zhang, D.L.; Liu, L.; Lee, M.R.; Wang, C.L.; Li, Z.; Mo, E.K.; Sung, C.K. (2011). Variabilities of two *Drechlerella dactyloides* isolates in Korea and high predacity against *Bursaphelenchus xylophilus*. – *Curr. Microbiol.* 62: 472–478.
- Witte, H.; Yang, J.; Borneman, J.; Smith-Becker, J.; Becker, J.O. – Assessment of sugar-beet fields for the presence of the nematophagous fungus *Dactylella oviparasitica* by a real-time PCR based strategy. Unpublished
- Wu, H.Y.; Kim, D.G.; Ryu, Y.H. – Unpublished
- Xu, L.L.; Yang, E.C.; An, Z.Q.; Liu, X.Z. – The Mesozoic Era was an important geological period for fungal evolution. Unpublished
- Yang, J.I.; Benecke, S.; Jeske, D.R.; Rocha, F.S.; Smith Becker, J.; Timper, P.; Becker, J.O.; Borneman, J. (2012). Population dynamics of *Dactylella oviparasitica* and *Heterodera schachtii*: toward a decision model for sugar beet planting. – *J. Nematol.* 44: 237–244.
- Yang, Y.; Liu, X.Z. (2005). *Dactylella coccinella* sp. nov., an anamorphic species. – *Mycotaxon* 91: 127–132.
- Yang, Y.; Yang, E.; An, Z.Q.; Liu, X.Z. (2007). Evolution of nematode-trapping cells of predatory fungi of the *Orbiliaceae* based on evidence from rRNA-encoding DNA and multiprotein sequences. – *Proc. Natl. Acad. Sci. USA* 104: 8379–8384.
- Yarwood, S.A.; Bottomley, P.J.; Myrold, D.D. (2010). Soil microbial communities associated with douglas-fir and red alder stands at high- and low-productivity forest sites in Oregon, USA. – *Microbial Ecology* 60: 606–617.
- Yin, B.; Valinsky, L.; Gao, X.; Becker, J.O.; Borneman, J. (2003). Identification of fungal rDNA associated with soil suppressiveness against *Heterodera schachtii* using oligonucleotide fingerprinting. – *Phytopathology* 93: 1006–1013.
- Yu, Z.F.; Kong, Y.J.; Zhang, Y.; Qiao, M.; Guo, J.W.; Zhang, K.Q. (2009). A new *Dactylella* species from *Orbilialba*. – *J. Microbiol.* 47: 265–269.
- Yu, Z.F.; Qiao, M.; Zhang, Y.; Qin, L.; Zhang, K.Q. (2011). *Pseudotriporiconidium*, a new anamorph genus connected to *Orbilialba*. – *Mycologia* 103: 164–173.
- Yu, Z.F.; Zhang, Y.; Qiao, M.; Zhang, K.Q. (2007). *Orbilialba dorsalis* sp. nov., the teleomorph of *Dactylella dorsalis* sp. nov. – *Cryptog. Mycol.* 28: 55–63.
- Zhang, Y.; He, X.X.; Baral, H.O.; Qiao, M.; Weiß, M.; Liu, B.; Zhang, K.Q.; Yu ZF (2015). *Orbilialba laevimarginata* sp. nov. and its asexual morph. – *Phytotaxa* 203: 245–253.
- Zhang, Y.; Qiao, M.; Weber, E.; Baral, H.O.; Hagedorn, G.; Zhang, K.Q.; Yu, Z.F. (2010a). *Arthrobotrys scaphoides* from China and Europe with a phylogenetic analysis including the type strain. – *Mycotaxon* 111: 291–300.
- Zhang, H.X.; Tan, J.L.; Wei, L.X.; Wang, Y.L.; Zhang, C.P.; Wu, D.K.; Zhu, C.Y.; Zhang, Y.; Zhang, K.Q.; Niu, X.M. (2012). Morphology regulatory metabolites from *Arthrobotrys oligospora*. – *J. Nat. Prod.* 75: 1419–1423.
- Zhang, Y.; Yu, Z.F.; Qiao, M.; Zhang, K.Q. (2006). A new species of *Orbilialba* from China. – *Cryptog. Mycol.* 27: 289–294.
- Zhang, Y.; Zhang, S.; Wang, M.; Bai, F.; Liu, X. (2010b). High diversity of the fungal community structure in naturally-occurring *Ophiocordyceps sinensis*. – *PLoS ONE* 5: E15570
- Zhang, Y.; Zhang, Y.R.; Dong, J.Y.; He, X.X.; Qiao, M.; Baral, H.O.; Zhang, K.Q.; Yu, Z.F. (2016). *Orbilialba tianmushanensis* sp. nov., a new member of the *O. luteorubella* group with an unusual asexual morph. – *J. Microbiol.* 54: 9–13.
- Zhang, Y.; Qiao, M.; Baral, H.O.; Xu, J.P.; Zhang, K.Q.; Yu, Z.F. (2020). Morphological and molecular characterization of *Orbilialba pseudopolybrocha* and *O. tonghaiensis*, two new species of *Orbiliaceae* from China. – *Int. J. Syst. Evol. Microbiol.* 70: 2664–2676. <https://doi.org/10.1099/ijsem.0.004088>
- Zhang, Y.R.; Zhang, Y.; Xu, J.P. – Molecular phylogeny and DNA barcoding of nematode-trapping hyphomycetes and related taxa. Unpublished
- Zheng, T.H.; Fang, W.X.; Cai, K.Z.; Wang, F.H.; Wang, B.B.; Luo, X.G.; Zhang, H.Y.; Zhao, M.W.; Wei, S.; Li, X.S. – Unpublished
- Zinger, L.; Shahnava, B.; Baptist, F.; Geremia, R.A.; Choler, P. (2009). Microbial diversity in alpine tundra soils correlates with snow cover dynamics. – *ISME Journal* 1–10.